

Exam : A00-240

Title: SAS Certified Statistical


Business Analyst Using SAS

9: Regression and Modeling

Credential

Version: DEMO

1.Refer to the ROC curve:



As you move along the curve, what changes?

- A. The priors in the population
- B. The true negative rate in the population
- C. The proportion of events in the training data
- D. The probability cutoff for scoring

Answer: D

- 2. When mean imputation is performed on data after the data is partitioned for honest assessment, what is the most appropriate method for handling the mean imputation?
- A. The sample means from the validation data set are applied to the training and test data sets.
- B. The sample means from the training data set are applied to the validation and test data sets.
- C. The sample means from the test data set are applied to the training and validation data sets.
- D. The sample means from each partition of the data are applied to their own partition.

Answer: B

- 3.An analyst generates a model using the LOGISTIC procedure. They are now interested in getting the sensitivity and specificity statistics on a validation data set for a variety of cutoff values. Which statement and option combination will generate these statistics?
- A. Score data=valid1 out=roc;
- B. Score data=valid1 outroc=roc;

- C. mode1 resp(event= '1') = gender region/outroc=roc;
- D. mode1 resp(event"1") = gender region/ out=roc;

Answer: B

- 4.In partitioning data for model assessment, which sampling methods are acceptable? (Choose two.)
- A. Simple random sampling without replacement
- B. Simple random sampling with replacement
- C. Stratified random sampling without replacement
- D. Sequential random sampling with replacement

Answer: A,C

5. Which SAS program will divide the original data set into 60% training and 40% validation data sets, stratified by county?

```
A proc surveyselect data=SASUSER.DATABASE samprate=0.6 out=sample;
      strata county;
   run;
B. proc sort data=SASUSER.DATABASE;
      by county;
   run;
   proc surveyselect data=SASUSER.DATABASE samprate=0.6 out=sample outall;
   run;
C. proc sort data=SASUSER.DATABASE;
      by county;
   run;
   proc surveyselect data=SASUSER.DATABASE samprate =0.6 out=sample outall;
      strata county;
   run;
D. proc sort data=SASUSER.DATABASE;
      by county;
   proc surveyselect data=SASUSER.DATABASE samprate = 0.6 out=sample;
      strata county;
   run;
```

- A. Option A
- B. Option B
- C. Option C
- D. Option D

Answer: C